
Name:		Off. Class:	Per:	Date:
Teacher:		Practice		Chemistry
is an isomer of $\begin{array}{c} H & H & H & Br \\ H & -C & -C & -C & -H^{2} \\ H & -L & + & H^{2} \\ H & -L & + & H^{2} \\ H & H & H & Br \end{array}$ A) Br H H H H H $\begin{array}{c} H & -C & -C & -C & -H \\ H & -C & -C & -C & -H \\ Br H & H & H & H \\ H & -C & -C & -H \\ H & H & H \\ H & H \\ H & H \\ H \end{array}$	B) H Br H H H $\stackrel{ }{}$ H Br H H H $\stackrel{ }{}$ H H H H $\stackrel{ }{}$ H H H H H H H D) H Br $\stackrel{ }{}$ H	properties. Both sub	th substances have rbon atoms, one ns. These two sulf each other each other ompound ydrocarbon compounds are in l2O4 24O10 nd CH3COOH and C2H5OH	oxygen atom, and six ostances must be
	 B) propane D) 1-chloropropane CH₃CH₂CH₂COOH? OH 		sure nt composition somer of the compoun 2COOH? COOH CH2COOH DH)CH2OH	-
A) $ \begin{array}{c} H & H & H & H \\ H & -C - C - C - C - O H \\ H & H & H & H \\ \end{array} $ C) $ \begin{array}{c} H & H & H & H \\ H & H & H & H \\ H & H &$	D) H O H H H-C-C-C-C-H H H H			


11. Base your answer to the following question on the information below.

The formula below represents a hydrocarbon.




In the space below, draw a structural formula for one isomer of this hydrocarbon.

12. Given the structural formula for butane:



Draw the structural formula of an isomer of butane.

13. Given the structural formula of pentane:



Draw a structural formula for an isomer of pentane.

14. Base your answer to the following question on the information below and on your knowledge of chemistry.

Diethyl ether is used as a laboratory and industrial solvent. The boiling point of diethyl ether at standard pressure is 34.6°C. The equation below represents a reaction that produces diethyl ether.

$$H \xrightarrow{H} C = C \xrightarrow{H} H \xrightarrow{H} C = C \xrightarrow{H} H \xrightarrow{H} H \xrightarrow{H} H^{2} O \xrightarrow{\text{catalyst}} H \xrightarrow{H} H \xrightarrow{H} H \xrightarrow{H} H \xrightarrow{H} H \xrightarrow{H} H^{2} O$$

Draw a structural formula for an isomer of the product that has the same functional group.

15. Base your answer to the following question on the information below and on your knowledge of chemistry.

Water, H₂O, and hexane, C₆H₁₄, are commonly used as laboratory solvents because they have different physical properties and are able to dissolve different types of solutes. Some physical properties of water and hexane are listed on the table below.


Solvent	Boiling Point (°C)	Melting Point (°C)	Vapor Pressure at 69°C (kPa)
H ₂ O	100.	0.	?
C_6H_{14}	69	-95	101.3

Physical Properties of H₂O and C₆H₁₄


Explain, in terms of molecular formulas and structural formulas, why 2,2-dimethylbutane is an isomer of hexane.

16. Base your answer to the following question on the information below and on your knowledge of chemistry.

The equation below represents a reaction between propene and hydrogen bromide.



Cyclopropane, an isomer of propene, has a boiling point of -33° C at standard pressure and is represented by the formula below.



Explain, in terms of molecular formulas and structural formulas, why cyclopropane is an isomer of propene.

17. Base your answer to the following question on the information below.

Many artificial flavorings are prepared using the type of organic reaction shown below.

Reactant 1Reactant 2Draw the structural formula of an isomer of reactant 2.

18. Base your answer to the following question on the information below.

Biodiesel is an alternative fuel for vehicles that use petroleum diesel. Biodiesel is produced by reacting vegetable oil with CH₃OH. Methyl palmitate, C₁₅H₃₁COOCH₃, a compound found in biodiesel, is made from soybean oil. One reaction of methyl palmitate with oxygen is represented by the balanced equation below.

 $2C_{15}H_{31}COOCH_3 + 49O_2 \rightarrow 34CO_2 + 34H_2O + energy$

Explain, in terms of both atoms and molecular structure, why there is no isomer of CH₃OH.

Base your answers to questions **19** and **20** on the information below and on your knowledge of chemistry.

There are several isomers of C_6H_{14} . The formulas and boiling points for two of these isomers are given in the table below.

Isomer	Formula	Boiling Point at 1 atm (°C)
1	H H H H H H 	68.7
2	$ \begin{array}{c} H \\ H \\ H \\ H \\ -C \\ -C$	49.7

19. Write the empirical formula for isomer 1.

20. Explain, in terms of intermolecular forces, why isomer 2 boils at a lower temperature than isomer 1.